DOI: 10.1002/ece3.10609

REVIEW ARTICLE

• WILEY

Density of wild honey bee, Apis mellifera, colonies worldwide

Oliver D. Visick 💿 | Francis L. W. Ratnieks

Laboratory of Apiculture and Social Insects (LASI), School of Life Sciences, University of Sussex, Brighton, UK

Correspondence

Oliver D. Visick, Laboratory of Apiculture and Social Insects (LASI), School of Life Sciences, University of Sussex, Brighton BN1 9QG, UK. Email: ov32@sussex.ac.uk

Funding information C.B. Dennis British Beekeepers' Research

Trust

Abstract

The western honey bee, Apis mellifera, lives worldwide in approximately 102 million managed hives but also wild throughout much of its native and introduced range. Despite the global importance of A. mellifera as a crop pollinator, wild colonies have received comparatively little attention in the scientific literature and basic information regarding their density and abundance is scattered. Here, we review 40 studies that have quantified wild colony density directly (n=33) or indirectly using genetic markers (n = 7) and analyse data from 41 locations worldwide to identify factors that influence wild colony density. We also compare the density of wild and managed colonies at a regional scale using data on managed colonies from the Food and Agriculture Organization (FAO). Wild colony densities varied from 0.1 to 24.2/km² and were significantly lower in Europe (average of 0.26/km²) than in Northern America (1.4/km²), Oceania (4.4/km²), Latin America (6.7/km²) and Africa (6.8/km²). Regional differences were not significant after controlling for both temperature and survey area, suggesting that cooler climates and larger survey areas may be responsible for the low densities reported in Europe. Managed colony densities were 2.2/km² in Asia, 1.2/km² in Europe, 0.2/km², in Northern America, 0.2/km² in Oceania, 0.5/km² in Latin America and 1/km² in Africa. Wild colony densities exceeded those of managed colonies in all regions except Europe and Asia. Overall, there were estimated to be between two and three times as many wild colonies as managed worldwide. More wild colony surveys, particularly in Asia and South America, are needed to assess the relative density of wild and managed colonies at smaller spatial scales.

KEYWORDS

Apis mellifera, beekeeping, biogeography, natural selection, population size, wild colony density

TAXONOMY CLASSIFICATION

Biogeography, Conservation ecology, Disease ecology, Entomology, Population ecology

1 | INTRODUCTION

Beekeeping with the western honey bee (*Apis mellifera* L.) dates back to ancient Egypt (Crane, 1999) and is now practiced on every continent except Antarctica, in both its native (Africa, Europe and Middle East) and introduced range (Americas, Asia and Oceania). In 2021, there were an estimated 102 million managed honey bee colonies worldwide (FAO, 2021), the majority being A. *mellifera*. A. *mellifera* pollinates approximately half of all globally important crops (Klein et al., 2007) and contributes over £100 billion to the global economy every

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

 $\ensuremath{\mathbb C}$ 2023 The Authors. Ecology and Evolution published by John Wiley & Sons Ltd.

year through pollination and honey production (Gallai et al., 2009). However, in addition to living in managed hives, A. *mellifera* colonies also live wild, typically nesting in cavities in trees (Figure 1) and buildings (Gambino et al., 1990; Saunders et al., 2021), but also in the ground and in rock crevices (Ratnieks et al., 1991) and occasionally not in a cavity (Boreham & Roubik, 1987; Saunders et al., 2021).

Apis mellifera is well-studied in the contexts of general biology and beekeeping, but wild colonies are less studied (Seeley, 2019). For example, the first comprehensive study of the characteristics of natural nests (Seeley & Morse, 1976) was not made until several decades after the discovery of the dance language (von Frisch, 1937). Wild colonies are sometimes (Thompson, 2012), and incorrectly, viewed as a by-product of beekeeping and have received comparatively little attention in the scientific literature (Kohl & Rutschmann, 2018). One important gap in our knowledge of *A. mellifera* is very basic: the density of wild colonies. Indeed, *A. mellifera* is listed as 'data deficient' on the ICUN Red List of Threatened Species (De la Rúa et al., 2014).

Wild A. *mellifera* colony densities were reviewed by Ratnieks et al. (1991) and found to range from 0.17 to 7.8/km² at 9 locations worldwide (Africa: 1, Europe: 1, Latin America: 3, North America: 4). Two decades later, using an indirect method based on population genetics, Jaffé et al. (2010) estimated the densities of both wild and managed colonies combined at 25 locations across its native range



FIGURE 1 Wild honey bee, *Apis mellifera*, colony in a tree cavity in southern England.

(Africa: 10, Asia: 2, Europe: 13), which ranged from 0.8 to 10.2/km². However, at most locations, the number of colonies detected using genetic markers could be accounted for by the estimated number of managed colonies, suggesting that wild colonies were rare or even absent in parts of its native range. However, indirect measures of colony density are subject to various limitations (reviewed by Utaipanon, Holmes, et al., 2019; Williamson et al., 2022), which make it difficult to detect wild colonies living at low densities in areas with many managed colonies (Kohl & Rutschmann, 2018).

Based on direct measures of wild colony density (Kohl & Rutschmann, 2018; Oleksa et al., 2013) and cavity density data, Requier et al. (2020) estimated the number of wild A. *mellifera* colonies that could be sustained in forests in a 4.6 million km² area of Europe. Forests made up 1.4 million km² (31%) of the study area and were estimated to contain approximately 80,000 wild colonies (0.057/km² of forest). This is only 2% of the number of managed colonies reported in this region by the FAO (Requier et al., 2020), although the estimated number of wild colonies was based on lower bound estimates of colony density and did not account for wild colonies present in habitats other than forests (Requier et al., 2020). Therefore, the actual number of wild colonies present in Europe is likely much higher and probably represents an important component of total colony numbers.

Here, we review all studies that have measured the density of wild A. mellifera colonies with a focus on studies that used direct surveys to count colonies in defined areas. In the 30 years since the review by Ratnieks et al. (1991), there has been an upsurge in interest in pollinators and the number of wild colony surveys reported in the literature has increased three-fold (see Section 4). A. mellifera is economically important and occurs worldwide and there are several reasons why information on the density of wild colonies is important, including the conservation of native subspecies (reviewed by Requier et al., 2019) and as sources of genetic variation in beekeeping (see below). Therefore, a comprehensive review is timely. In addition, we further analyse the data set to investigate factors that influence wild A. mellifera colony density. We also compare the density of wild and managed colonies at a regional scale using data on managed colonies from the FAO and make a tentative estimate for the number of wild A. mellifera colonies worldwide.

2 | WHY IS WILD A. MELLIFERA COLONY DENSITY IMPORTANT?

Information on the density of wild A. *mellifera* colonies, in both its native and introduced ranges, is of both ecological and economic importance. For instance, measures of wild colony density in agricultural areas, in combination with information on foraging distances (Couvillon et al., 2014), could be used to assess their potential contribution to crop pollination. Indirect measures of wild colony density have been lower in agricultural areas compared to natural habitats (Hinson et al., 2015; Jaffé et al., 2010), possibly due to a lack of suitable nesting sites in the former (Oleksa et al., 2013). Therefore, the

WILEY

contribution of wild colonies to crop pollination in large-scale farming operations is probably low compared to managed colonies which can be placed at very high densities for short periods of time, such as 2 per acre ($500/km^2$) for almond pollination (Mader et al., 2010) and 6.7 per acre ($1650/km^2$) for onion pollination (literature average, Delaplane & Mayer, 2000).

Measures of colony density can also be used to address concerns regarding the impact that both managed and wild colonies of *A. mellifera* have in their introduced range. Proposed threats include competition with native pollinators for floral resources (Paini, 2004), competition with native vertebrates for nest sites (Oldroyd et al., 1994; Pacífico et al., 2020), pollination of exotic weeds (Goulson & Derwent, 2004) and the introduction of exotic pests and disease (Taylor et al., 2007). In Australia, measures of colony density have been used to determine whether wild colonies pose a significant threat to native ecosystems and, in extreme cases, whether eradication would be possible (Hinson et al., 2015; Oldroyd, 1998; Oldroyd et al., 1994).

Repeated surveys can also be used to monitor changes in wild colony density over time. For example, in the Arnot Forest in western New York State, three surveys spanning over 30 years (1978, 2002 and 2011) showed that the introduction of the ectoparasitic mite, *Varroa destructor*, in the 1990s had no long-term effect on the density of wild A. *mellifera* colonies living in the area (Seeley, 2007; Seeley et al., 2015; Visscher & Seeley, 1982) even though *Varroa* became established between the first and second surveys. Similarly, in The Welder Wildlife Refuge, Texas, two surveys conducted both during (1991–2001) and after (2013) the invasion of Africanised bees showed that wild colony densities remained high over a decade after the invasion (5.4/km²) and that wild colonies retained a stable mixture of European- and Africanderived genetics. These studies are important in showing that wild populations of *A. mellifera* can remain stable over long periods despite significant changes in the bees themselves or their pests.

Importantly, the role of natural selection on colony survival will be affected by the number of wild colonies present in an area. Many types of disease management, including the use of chemicals to kill Varroa mites, play an important role in the survival of managed colonies (van Dooremalen et al., 2012), but will likely reduce the effect of natural selection for disease resistance (Neumann & Blacquiere, 2016). Several wild and unmanaged populations of A. mellifera have been shown to possess natural mechanisms that reduce Varroa population growth (Mondet et al., 2020) including short post-capping durations (Le Conte et al., 2007; Oddie et al., 2018), cell recapping (Hawkins & Martin, 2021; Oddie & Dahle, 2021) and Varroa-sensitive hygiene (Harris, 2007; Panziera et al., 2017). In a typical population of A. mellifera, where wild and managed colonies can interbreed over long distances (De la Rúa et al., 2013), heritable traits that confer long-term resistance to Varroa will likely increase in frequency more rapidly when a greater proportion of colonies are wild and exposed to natural selection (Requier et al., 2019; Youngsteadt et al., 2015). These traits are also present in managed colonies at low frequencies and can be increased via artificial selection and queen rearing (Bigio et al., 2014; Büchler et al., 2010; Pérez-Sato et al., 2009; Rinderer et al., 2010).

3 | METHODS

3.1 | Choice of studies

Here, we review 55 reports of wild honey bee colony density from a total of 40 studies published between 1971 and 2022, although the actual data from Galton (1971) originates from the late 1600s (Seeley, 2019). 33 reports come from regions where *A. mellifera* is introduced (Australia: 16, Latin America: 7, USA: 10), versus 22 from within *A. mellifera*'s native range (Africa: 13, Europe: 9). Interestingly, there have been no reports of established wild populations of *A. mellifera* in eastern or southern Asia despite their increasing use in commercial beekeeping. This is suggested to be due to the parasites and diseases of other species of honey bee present in these regions (e.g. *Apis cerana*; Oldroyd & Nanork, 2009; see Section 5).

Direct measures of wild colony density (n=35, Table 1) were made using one or more of the following methods. Direct searches were the most common (n=19) and involved looking for bees going in and out of nesting cavities. Nine surveys were made using bee-lining, which involves following marked bees back to their nest by recording the direction they depart from a food source and the time it takes them to return (Seeley, 2016). Four surveys were made using local knowledge, which involved communications with local residents, landowners and African honey hunters (Kajobe & Roubik, 2006; Schneider & Blyther, 1988). Four surveys were made using data from either forest beekeeping in Russia (n=1), in which honey from wild colonies living in trees is harvested (Seeley, 2019) or colony removal records from urban areas (n=3).

Indirect measures of wild colony density (n=19, Table 2) were made by taking samples of honey bees from either the worker progeny of gueens mated in an area of interest (Arundel et al., 2014; Jaffé et al., 2010; Moritz et al., 2007) or, more commonly, from drones trapped at drone congregation areas (DCAs; Arundel et al., 2013; Hinson et al., 2015; Jaffé et al., 2010; Moritz et al., 2007, 2008, 2013; Utaipanon, Holmes, et al., 2019; Utaipanon, Holmes, et al., 2021; Utaipanon, Schaerf, et al., 2021). The number of colonies within a given radius of the mating apiary or DCA is then inferred by the number of unique genotypes present in the sample of worker progeny or trapped drones, respectively. Indirect measures of wild colony density were only included if they excluded the effect of managed colonies on the total number of colonies detected. For example, Jaffé et al. (2010), Arundel et al. (2014), Hinson et al. (2015) and Utaipanon, Schaerf, and Oldroyd (2019) sampled sites with little or no managed colonies within drone flight distance.

3.2 | Area surveyed

The area over which wild colonies were located (direct) or detected (indirect) was, in most cases, explicitly stated by the authors in their calculation of wild colony density. Survey areas were sometimes given as the total area of multiple plots of a standard size ($<0.05 \text{ km}^2$ each; Darchen, 1972; Goodman & Hepworth, 2004;

					982)	1987)	- (1988)						۵ Oliver, D	ir (1996) ^h		rth (2004)	rth (2004)		(900					
Reference	Galton (<mark>1971</mark>) ^a	Kerr (1971) ^a	Darchen (1972) ^d	Taber (1979)	Visscher and Seeley (1	Boreham and Roubik (Schneider and Blythe	Wenner (1989)	Morse et al. (1990)	Ratnieks et al. (1991)	Oldroyd et al. (1994)	Danka et al. (1994)	Paton, DC, Jansen, L { (unpublished) ^g	McNally and Schneide	Oldroyd et al. (1997)	Goodman and Hepwo	Goodman and Hepwo	Baum et al. (2005)	Kajobe and Roubik (2	Seeley (<mark>2007</mark>)	Baum et al. (2008)	Oleksa et al. (2013)	Seeley et al. (2015)	
Colonies/ km ²	0.17-0.96	4.4	10	2.9-5.1	1.06	1.2-2.1	7.8	0.25	2.3	5-9	77.1	3.02	0.11-40	4.2	40-148	2.9	92	1.6-12.64	12	0.94	0.35-1.12	0.1	1.06	
Survey area (km²)	10-88	1.80	0.30 ^c	3.14	8.5	50	6	230	4.2	1-2.1	0.35 ^c	12.6	0.09-20	19.3	0.25	0.35 ^c	0.91 ^c	6.25	1.74	8.5	924.1 ^k	458.1	8.5	
No. colonies	3-50	8	ю	9-16	6	60-105 ^e	47	58 ^f	11	5-13	27	38	1-60	81	10-37	4	84	10-79 ⁱ	20	ω	323-1035	45	6	
No. plots/ sites	4	15	94	I	Т	Т	Т	I	I	ო	7	I	Ŋ	I	5	35	30	I	87	I	I	170	I	
Main cavity type	Tree hollows	Tree hollows	Tree hollows	Rocks crevices	Trees hollows	Man-made	Trees hollows	Rocks crevices	Man-made	Tree hollows	Tree hollows	Tree hollows	I	Tree hollows	Tree hollows	Tree hollows	Tree hollows	Tree hollows	Tree hollows	Tree hollows	Man-made	Tree hollows	Tree hollows	
Survey method(s)	Colony records	Direct search ^b	Direct search ^b	Direct search	Bee-lining	Colony records	Local Knowledge	Bee-lining	Local knowledge	Search & local	Direct search	Bee-lining, search & local	Direct search ^b	Search & local	Direct search	Direct search	Direct search	Direct search	Local knowledge ^j	Bee-lining	Colony records	Direct search	Bee-lining	
Habitat	Temperate forest	Woodland	Dry forest	Semi-desert canyon	Temperate forest	Mixed	Semi-desert	Arid Island	Urban/suburban	Agricultural land	Riparian woodland	Patchy tropical forest	Conservation parks	Semi-desert	Riparian woodland	Box-ironbark Forest	Riparian woodland	Coastal prairie	Montane forest	Temperate forest	Urban/suburban	Rural avenues	Temperate forest	
Longitude	44.0*	-52.4*	-5.3*	-112.0	-76.7	-79.7*	23.4	-119.7	-76.5	-92.3	141.9	-85.6	139.1*	23.4	141.9	146.6*	145.4*	-97.4	29.7	-76.7	-111.0	19.9*	-76.7	
Latitude	56.3*	-18.5*	5.6*	34.7	42.3	9.1*	-19.6	34.0	43.5	14.9	-35.6	10.4	-35.6*	-19.6	-35.6	-36.1*	-36.4*	28.1	-0.6	42.3	32.2	53.7*	42.3	
Location	Morozov Estate	Goiás & Mato Grosso	Lamto Savannah	Cottonwood, AZ	Arnot Forest, NY	Panama Canal	- Okavango	Santa Cruz Island, CA	Oswego, NY	Tapachula	Wyperfeld NP, Vic	Belén	South Australia	Okavango	Wyperfeld NP, Vic	Box-ironbark Forest, Vic	Goulburn Valley, Vic	Welder Wildlife Refuge, TX	Bwindi Impenetrable NP	Arnot Forest, NY	Tucson, AZ	Northern Poland	Arnot Forest, NY	
Country	Russia (N)	Brazil (I)	lvory Coast (N)	USA (I)	USA (I)	Panama (I)	Botswana (N)	USA (I)	USA (I)	Mexico (I)	Australia (I)	Costa Rica (I)	Australia (I)	Botswana (N)	Australia (I)	Australia (I)	Australia (I)	USA (I)	Uganda (N)	USA (I)	USA (I)	Poland (N)	(I) VSN	

TABLE 1 Thirty-six reports from 33 studies that have quantified wild A. mellifera colony density directly.

\sim	-
7	5
Ĩ	5
. <u>-</u>	
t	
Ē	2
<u> </u>	2
~	1
-	1
Г Т 1	
BIF 1	1
ABIF 1	1

Country	Location	Latitude	Longitude	Habitat	Survey method(s)	Main cavity type	No. plots/ sites	No. colonies	Survey area (km ²)	Colonies/ km ²	Reference
South Africa (N)	Cape Point	-34.4*	18.5*	Fynbos (heathland)	Bee-lining & search	Rock crevices	I	59	50.4 ^m	1.17	Tribe et al. (2017)
Germany (N)	Hainich NP	51.1	10.4	Temperate forest	Bee-lining	Tree hollows	T	4-6 ⁿ	30.6-43.2	0.13	Kohl and Rutschmann (2018)
Germany (N)	Swabian Alb	48.4	9.5	Temperate forest	Direct search ^o	Tree hollows	I	7	61.3	0.11	Kohl and Rutschmann (2018)
(I) VSN	Shindagin Hollow	42.3	-76.3	Temperate forest	Bee-lining	Tree hollows	I	5	5.18	0.97	Seeley and Radcliffe (2018)
Zambia (N)	Lusaka	-14.2	30.2	Miombo/mopane	Direct search	Tree hollows	2	29	1.2	24.2	Coppinger et al. (2019)
Scotland (N)	Cawdor Wood	57.5	-3.9	Woodland	Bee-lining	Tree hollows	T	4	31.1	0.13	Seeley and Chilcott (2020)
Serbia (N)	Belgrade	44.8	20.5	Urban/suburban	Colony records	Man-made	I	460 ^p	224	2.05	Bila Dubaić et al. (2021)
Spain (N)	Xinzo de Limia	42.1	-7.7	Agricultural land	Direct search ^q	Man-made	ı	23-29	136	0.17-0.22	Rutschmann et al. (2022)
Australia (I)	Albury, NSW	-35.5*	147.8*	Patchy woodland	Direct search ^r	Man-made	10	36	0.53 ^c	68.6	Cunningham et al. (2022)
Germany (N)	Coburg & Lichtenfels	50.2*	11^*	Temperate forest	Direct search ^o	Tree hollows	I	0-21	22.1-59.1	0-0.36	Kohl et al. (2022)
Germany (N)	Swabian Alb	48.4	9.5	Temperate forest	Direct search ^o	Tree hollows	I	0-20	48.6-116.9	0-0.21	Kohl et al. (2022)
Australia (I)	Adelaide, SA	-35.0	138.6	Arboretum/urban	Direct search	Tree hollows	Т	31-81	1.34-7.65	23.1-10.59	Williamson et al. (2022)
<i>lote</i> : The letter	in column 1 corresponc	ls with whe	ther each cou	ntry is in A. mellifera's	native (N) or introdu	Iced (I) range. A	dash (–) ir	n column 8 ir	idicates that a	a contiguous a	area was surveyed. An asterisk (*)

next to a coordinate in column 3 or 4 indicates that it is was either approximated or averaged between sites. ^aFrom Ratnieks et al. (**1991**).

^bSurvey method not reported (assumed to be direct search).

^cDensities measured in areas $< 1 \, \text{km}^2$ were included in the table for completeness but omitted from analyses.

^dFrom Kajobe and Roubik (2006).

^eRange of colonies removed over 4-year period. Ratnieks et al. (1991) reported both colonies and swarms.

 $^{
m f}$ Exact number of colonies not reported. Calculated from colony density and area surveyed.

^gFrom Paton (1996).

^hCombined with data from Schneider and Blyther (1988).

ⁱFrom Rangel et al. (2016).

 $^{\rm j}$ Colonies located by guides and indigenous honey hunters. $^{\rm k}$ Survey area calculated from reported colony density and numbers.

¹Forty-five colonies located on 142 km of rural avenues. Colony density estimated via avenue density.

^mSearched 65% of the Cape Point section of the Table Mountain National Park (77.5 km²).

"Nest not actually located (location inferred by bee-lining).

^oOnly searched tree hollows that were previously occupied by black woodpeckers.

^pFour hundred sixty colonies identified during 7-year study period. Only c. 20% were continuously occupied during study period.

^qOnly searched concrete power poles.

^rOnly searched nest boxes.

WILEY

ŝ
er
¥
na
J
it:
Ű
80
60
Sil.
Ë
<u>≥</u>
5
.≝
pu
.=
÷
ů.
de
2
Ы
0
5
БГ
ΠĘ
Jel
2
A
lld
Ī
p
fie
÷
lar
Ъ
ē
٩٧
t
Ja
÷
es
ipi
ŝt
Ľ,
Š
Ś
Ē
õ
f f
rts
ō
je,
n r
ē
et
Ĩ.
Ζ
~
-
Ш.
Ω
<
H .

						1					
Country	Location	Latitude	Longitude	Habitat	Sampling method	Drones genotyped	No. N sites co	o. olonies	Survey area (km²)	Colonies/km ²	Reference
South Africa (N)	Gauteng ^a	-25.9	28.6	Nature reserve	Drone trapping	96-127	3.	2-44	2.5	12.8-17.6	Moritz et al. (2007)
South Africa (N)	Tswalu game reserve ^a	-27.2	22.4	Nature reserve	Drone trapping	49	- 2(5.5	2.5	10.6	Moritz et al. (2008)
South Africa (N)	Gauteng ^a	-25.9	28.6	Nature reserve	Drone trapping	96-191	3 2;	3-37	2.5	9.2-14.8	Jaffé et al. (2010)
South Africa (N)	Tswalu game reserve ^a	-27.2	22.4	Nature reserve	Drone trapping	148	- 2	6	2.5	11.6	Jaffé et al. (2010)
South Africa (N)	Jonkershoek ^a	-34.0	18.9	Nature reserve	Drone trapping	96	- 4(0	2.5	16	Jaffé et al. (2010)
South Africa (N)	Pietermaritzburg ^a	-29.6	30.5	Nature reserve	Drone trapping	96	- 5,	~	2.5	20.8	Jaffé et al. (2010)
Sudan (N)	AI-faw ^a	14.2	34.3	Agricultural land	Worker progeny	I	- 2	m	4.5	5.1	Jaffé et al. (2010)
Mexico (I)	Chiapas N. ^b	16.7*	-92.4*	Agricultural land	Drone trapping	89-96	5 33	8-52	2.5	20.8-15.2	Moritz et al. (2013)
Mexico (I)	Chiapas S. ^b	14.9*	-92.5*	Mangrove & mango plantation	Drone trapping	89-183	й ч	4-43	2.5	13.6-17.2	Moritz et al. (2013)
Mexico (I)	Yucatan ^b	21.1^{*}	-88.9*	Deciduous forest	Drone trapping	92	2 3.	7-42	2.5	14.8-16.8	Moritz et al. (2013)
Australia (I)	Wyperfeld NP & Yaapeet, Vic ^c	-35.6*	142.0*	Mallee ^e	Worker progeny	I	2	5-29	4.5 ^f	5.6-6.4	Arundel et al. (2014)
Australia (I)	Puckapunyal N. & S., Vic ^c	-36.96*	144.9*	Box ironbark forest	Worker progeny	I	2 3 [.]	-11 ,	4.5 ^f	0.67-2.4	Arundel et al. (2014)
Australia (I)	Puckapunyal E. & Dookie, Vic ^c	-36.6*	145.4*	Box ironbark forest ^e	Worker progeny	I	2 14	4-18	4.5 ^f	3.1-4	Arundel et al. (2014)
Australia (I)	Marysville & Eildon, Vic	-37.4*	145.8*	Sclerophyll forest ^e	Worker progeny	I	2 1:	3-15	4.5 ^f	2.9–3.3	Arundel et al. (2014)
Australia (I)	Barrington Tops, NSW ^d	-32.2*	151.7*	Sclerophyll forest ^e	Drone trapping	70-123	4 14	4-23.5	2.5 ⁸	5.6-9.4	Hinson et al. (2015)
Australia (I)	Weddin Shire, NSW ^d	-33.6*	148.1^{*}	Woodland ^e	Drone trapping	14-241	4 6	-49	2.5 ⁸	2.4-19.6	Hinson et al. (2015)
Australia (I)	Wimmera, Vic ^d	-35.6*	142.1^{*}	Mallee ^e	Drone trapping	62-278	4 18	3-37.5	2.5 ⁸	7.2-15	Hinson et al. (2015)
Australia (I)	Grong Grong, NSW	-34.8	146.7	Mixed farming	Drone trapping	24-251	15 23	36	86.5 ^h	2.73	Utaipanon, Schaerf, and Oldroyd (2019)
Australia (I)	Currawarna, NSW	-35.0	147.1	Mixed farming	Drone trapping	22-216	15 20	55	86.5 ^h	3.06	Utaipanon, Schaerf, and Oldroyd (2019)
	-	1 • •	•		- - -	•		-		•	

Note: All indirect measures have been made over the last 15 years and are mainly limited to the Southern hemisphere. The letter in column 1 corresponds with whether each country is in A. mellifera's native (N) or introduced (I) range. An asterisk (*) next to a coordinate in column 3 or 4 indicates that it is was either approximated or averaged between sites.

 $^{\rm a}{\rm No}$ managed colonies within a 2.5–3 km radius of the sampling site.

^b No difference in number of colonies detected in areas with high and low beekeeping.

^cBeekeeping not practiced in study location for 40–50years.

 $^{\mathsf{d}}\mathsf{B}\mathsf{e}\mathsf{e}\mathsf{k}\mathsf{e}\mathsf{e}\mathsf{p}\mathsf{i}\mathsf{n}\mathsf{g}$ either prohibited from study locations or absent during time of sampling.

^eSurvey location consisted of paired disturbed and undisturbed sites.

^fOriginal survey area and density calculated using agent-based model from Arundel et al. (2012).

^gOriginal survey area and density calculated using agent-based model from Arundel et al. (2013).

^hSurvey area calculated at two 7km transects with typical drone flight range of 3.75km.

Kajobe & Roubik, 2006; Oldroyd et al., 1994, 1997) or the area of a circle with a given radius (Danka et al., 1994; Morse et al., 1990; Seeley & Radcliffe, 2018; Seeley & Chilcott, 2020; Taber, 1979), but in most cases the origin of survey areas was not given. In cases where survey areas were not explicitly stated, they were estimated by either dividing the number of colonies by the reported density (Baum et al., 2008; McNally & Schneider, 1996) or by using other information provided by the authors. For example, survey areas in Oleksa et al. (2013) and Kohl et al. (2022) were calculated using the density of rural avenues and cavity trees, respectively.

Areas surveyed using indirect measures were assumed to be either 2.5 km² (drone trapping) or 4.5 km² (worker progeny), based on the assumption that drones mate at a median distance of 900m from their colony (Taylor & Rowell, 1988) and that queens mate over an area approximately 1.8 times as large (Jaffé et al., 2010). An exception to this is Utaipanon, Schaerf, and Oldroyd (2019), who measured their own drone mating distances and found that drones caught along two 7-km transects in New South Wales were sampled from a much larger area of 86.5 km². Honey bee mating distances can vary significantly (Jensen et al., 2005) and even a small increase in distance can have a large effect on the resulting area (Utaipanon, Schaerf, & Oldroyd, 2019).

3.3 | How multiple surveys at a location were combined

Data from each survey location were combined to produce a single value for wild colony density. In some locations, data were combined over multiple years (Baum et al., 2005, 2008; Bila Dubaić et al., 2021; Boreham & Roubik, 1987; Kohl et al., 2022; Paton, 1996; Rutschmann et al., 2022; Taber, 1979) or across multiple sites or plots within a wider general location (Darchen, 1972; Galton, 1971; Goodman & Hepworth, 2004; Ilyasov et al., 2015; Kajobe & Roubik, 2006; Kerr, 1971; Oldroyd et al., 1994, 1997; Oleksa et al., 2013; Paton, 1996;

Ratnieks et al., 1991). In the former, the mean number of wild colonies located each year was divided by the survey area, which was assumed to remain constant, as some colonies reported in each year would have been the same colonies as the previous year. For instance, using colony removal records, Baum et al. (2008) inferred the location of wild colonies in a 900km² area of Tucson, Arizona, during the invasion of Africanised bees from 1994 to 2001. The mean number of colonies located each year was 644.7, which produced a combined density of 0.7/km². In cases where survey area did not remain constant (Kohl et al., 2022; Williamson et al., 2022), only data from the most recent survey were used to calculate density.

To combine data collected at different sites or plots within a survey location the total number of colonies located (direct) or detected (indirect) was divided by the total survey area. For example, Ratnieks et al. (1991) located 27 colonies in three sites near Tapachula, Mexico, with a total area of 4.1 km^2 , which resulted in a combined density of $6.6/\text{km}^2$. The same method was used to calculate regional densities of managed colonies using the total landmass (km²) of countries where FAO data are available (n = 117).

3.4 | Criteria that excluded a study

In cases where a location had been surveyed by different studies using the same survey method, only data from the most recent study were included in the analysis. This includes the Arnot Forest, Okavango, Welder Wildlife Refuge, Wyperfeld National Park (Wimmera), Gauteng and Tswalu Game Reserve (Figure 2). Studies that surveyed a total area of $<1 \,\mathrm{km^2}$ (Cunningham et al., 2022; Darchen, 1972; Goodman & Hepworth, 2004; Oldroyd et al., 1994, 1997) were not included in the analysis because these often produced unrepresentative high densities (>50/km², Table 1) that were probably not representative of the surrounding habitats.



FIGURE 2 Forty-one locations worldwide where wild *Apis mellifera* colony density has been quantified directly (red) or indirectly via genetic markers (blue). The grey dashed line indicates the approximate northern limit of wild colonies in Europe (60 degrees), which is based on the distribution of lime and hazel trees (Requier et al., 2019). Countries in green (n = 117) have FAO data regarding the number of managed honey bee colonies. Map generated in QGIS (v3.16.11).

3.5 | Statistical analyses

All analyses were performed in R (v4.3.1; R Core Team, 2023) and all plots were made using ggplot2 (Wickham, 2016). We analysed data from a total of 41 locations where wild colony density had been quantified either directly (n=25) or indirectly using genetic markers (n=16; Table S1). Each survey location was allocated to one of five broad geographical regions (Africa, Europe, Latin America, Northern America, Oceania). The Americas were split into Northern (USA and Canada) and Latin America rather than North and South America because only one report of wild colony density has been made in South America (Kerr, 1971). Each location was also allocated to one of three categories of land use (disturbed, undisturbed, mixed) based on the level of human disturbance (similar to Arundel et al., 2014; Hinson et al., 2015). For instance, natural habitats, such as nature reserves and unmanaged woodland, were listed as undisturbed (n=20), whereas agricultural land and urban areas were listed as disturbed (n = 12) and locations consisting of both were listed as mixed (n=9; see Tables 1 and 2 for a full list of habitats).

Climate data were obtained in the form of raster datasets with a global coverage. The mean value of each variable was extracted from a 50km radius around each survey location using the Zonal Statistics tool in QGIS (v3.16.11). Simulated monthly temperature (°C) and precipitation (mm) data, with a spatial resolution of 0.5 by 0.5 degrees, were obtained from the Climatic Research Unit Time-Series (v4.04; Harris et al., 2020) and averaged from 1970 to 2020. Monthly net primary productivity (gC/m³/day) data, with a spatial resolution of 0.1 by 0.1 degrees, were obtained from NASA's terra MODIS satellite (product key: MOD17, v6.1; Running & Zhao, 2021) and averaged from 2001 to 2015.

3.6 | Model selection

Generalised linear models (GLMs) with Gamma error distributions were used to test for the effect of region, land use, climate and survey area on density at the 41 locations. Models with and without a variable were compared using their Akaike information criterion (AIC), a measure of goodness of fit that penalises models with more variables. A lower AIC indicates that a model better fits the data, although a difference in AIC (Δ AIC) of <2 is considered non-significant. Models with only one variable were compared with the null model, which only includes an intercept term. Tukey HSD tests were used to test for regional differences in density after controlling for other variables such as climate and survey area. Tukey tests were run using *multcomp* (Hothorn et al., 2008) and *p* values were adjusted using the Bonferroni method. Geometric means are given for density and area because they both occur on a logarithmic scale.

3.7 | FAO data on managed colonies

FAO data on managed colonies were available for 117 countries (Africa: 24, Asia: 26, Europe: 33, Latin America: 22, Oceania: 10,

Northern America: 2; Table S2; Figure 2). Only the most recent reports of managed colony numbers were used to calculate density. For most countries, these were made in 2021, except for Guadeloupe (1990), the Netherlands (1987), the United Kingdom (1987) and Belgium (2017). The FAO does not specify the species of honey bee that are managed in each country, and in particular, the proportion of managed colonies in southern and eastern Asia that are *A. cerana*. However, it is believed that, even in eastern Asia, most managed colonies are *A. mellifera* (Osterman et al., 2021). *A. cerana* is native only to Asia, but not including western Asia and had been introduced into New Guinea and Queensland, Australia (Koetz, 2013).

4 | RESULTS

4.1 | Regional variation in wild colony density

Wild colony densities reported in the literature were highly variable, ranging from 0.1/km² in Northern Poland to 148/km² in Australia (Table 1). Densities in our sample of 41 locations worldwide (Figure 2), ranged from 0.1 to 24.2/km² and fit a Gamma distribution with a geometric mean of 2.5/km². Region had a significant effect on density (Δ AIC=30.7). Densities reported in Europe (average of 0.26/km²) were significantly lower than Northern America (1.4/km², *p*=.01), Oceania (4.4/km², *p*<.001), Latin America (6.7/km², *p*<.001) and Africa (8.4/km², *p*<.001; Figure 3). Densities reported in Northern America (*p*=.022) and Latin America (*p*=.033) but not Oceania (*p*=.39). Land use had no effect on density (Δ AIC=-2.2).

4.2 | Effect of climate

There was a significant positive correlation between wild colony density and mean annual temperature ($\Delta AIC = 18.2$). Temperature also had a significant quadratic component ($\Delta AIC = 9.9$) with densities starting to decrease at mean annual temperatures exceeding 23°C (Figure 4). This model was not a better fit when other variables were included, such as mean monthly rainfall ($\Delta AIC = -1.5$) and net primary productivity ($\Delta AIC = 1.6$), although the latter was borderline significant. Region still had a significant effect on density after controlling for temperature ($\Delta AIC = 5.6$), with densities remaining significantly lower in Europe (p < .05) but not Northern America (p > .1), compared to the other three regions.

4.3 | Effect of survey area

Survey area varied considerably over nearly 3 orders of magnitude, even after excluding studies that surveyed <1 km (see methods), ranging from 1.2 km² in Zambia to 924 km² in Tucson, Arizona. Survey area had a significant negative correlation with density (Δ AIC=24.3; Figure 4) and was significantly larger in Europe (average of 111 km²)

NILEY



FIGURE 4 Effect of mean annual temperature (left) and survey area (right) on wild Apis mellifera colony density. Temperature had a significant quadratic component with densities decreasing at mean annual temperatures exceeding $23^{\circ}C$ ($R^2 = .55$). Density had a negative correlation with survey area ($R^2 = .53$). Fit lines and 95% confidence intervals were calculated using a linear model and log-transformed densities and areas.

and Northern America (18 km²), compared to Latin America (8.7 km², p < .01) and Africa (4.7 km², p < .01) but not Oceania (16.2 km², p > .1). Region still had a significant effect on density after controlling for survey area (Δ AIC = 13.2). However, region no longer affected density after controlling for both survey area and mean annual temperature (Δ AIC = -1.2).

4.4 | FAO managed colony numbers

Using recent data from the FAO, the number and density of managed colonies were calculated to be 1.4 million (0.18/km²) in Oceania, 3.4 million (0.18/km²) in Northern America, 8.2 million (0.47/km²) in Latin America, 18.2 million in Africa (0.96/km²), 25.4 million (1.2/km²)

in Europe and 45.3 million (2.2/km²) in Asia. Managed colony densities were lower than mean wild colony densities in all regions except Europe and Asia (Figure 5).

5 | DISCUSSION

5.1 | Regional variation in wild colony density

Our results show that wild A. *mellifera* colony densities vary over approximately 2.4 orders of magnitude from 0.1 to 24.2/km² with a mean of 2.5/km². Wild colony densities were highest in sub-Saharan Africa with an average of 6.8/km². African A. *mellifera* swarm frequently and maintain smaller colonies (McNally & Schneider, 1996),

10 of 15



FIGURE 5 Density of wild and managed A. mellifera colonies. Wild colony densities are the mean densities plotted in Figure 3. Managed colony densities were calculated using the most recent data from the FAO (2021) and the landmass of countries where data are available (n = 117). Wild colony densities exceed those of managed colonies in all regions except Europe. There have been no reports of wild colony density in Asia.

which enables them to utilise a wider range of nesting cavities (Baum et al., 2005) and can even build nests in the open (Saunders et al., 2021). These life history traits will likely have a large effect on the spatial distribution of wild colonies in areas where the maiority are of African descent. For instance, in the southern USA, wild Africanised honey bees (hybrids of A. mellifera scutellata and European subspecies) were more likely to occupy man-made nesting cavities (Baum et al., 2008) and form colony aggregations (Baum et al., 2005) than European colonies living in the same area.

The high wild colony densities reported in Africa are also in keeping with the fact that honey and brood are commonly harvested from wild colonies by tribes in Central Africa (Crittenden, 2011; Kajobe & Roubik, 2006). Traditional beekeeping in Africa also relies on the colonisation of log hives by wild swarms (Gratzer et al., 2021). In addition, the greater honeyguide (Indicator indicator) has evolved a sophisticated mutualism with humans in the detection and predation of wild colonies (Spottiswoode et al., 2016; Wood et al., 2014). Wild honey bees are clearly an important part of human life in Africa and it is unlikely that these complex relationships would have evolved if wild colonies were rare.

Wild colony densities in Europe (average of $0.26/\text{km}^2$) were significantly lower than all other regions and were approximately 25 times lower than those reported in Latin America and Africa (Figure 3). These results probably reflect, in part, differences in the carrying capacity of temperate and tropical ecosystems. For instance, it is possible that temperate ecosystems do not produce enough nectar or pollen to support many wild colonies and that they naturally occur at lower densities in Europe (Kohl & Rutschmann, 2018). In a sub-tropical prairie, Baum (2003) found that the nectar and pollen produced by plants within a 6.3 km² study area could support between 407 and 3161 wild colonies, but the number of colonies located during an 11-year survey rarely exceed 75 (12/km²; Baum et al., 2005), suggesting that wild colony density was limited by other factors. Here, we show that wild

colony densities remained significantly lower in Europe, but not Northern America, after controlling for climate.

It is possible that human activities are partly responsible for the low densities reported in Europe. For instance, it has been suggested that natural nest sites in old trees are lacking in many parts of Europe due to historical changes in land use (Carreck, 2008; Kohl et al., 2023; Kohl & Rutschmann, 2018; Oleksa et al., 2013) and that wild colonies are affected by the high density of managed colonies in this region through competition for food and the transfer of exotic pests and maladaptive genetics (Requier et al., 2019). However, estimates from 3 national parks in the Southern Urals (Ilvasov et al., 2015) suggested that wild colonies are living in Russian bee forests at similar densities (0.3/km²) to those recorded in the late 1600s (Galton, 1971), prior to major land use change (Chorley, 1981) and the advent of modern beekeeping in Europe (Crane, 1999). Although the historical density of wild colonies is not known for other parts of Europe.

It is also possible that wild colony densities were underestimated in parts of Europe. Four European studies inferred densities over large areas (average of 111 km²) by only searching for wild colonies in a specific habitat or nesting site, such as trees on rural avenues (Oleksa et al., 2013), black woodpecker (Dryocopus martius) nests in forests (Kohl & Rutschmann, 2018; Kohl et al., 2022) and concrete 'power poles' in an agricultural landscape (Rutschmann et al., 2022). Therefore, density was probably underestimated because wild colonies present in other habitats or nest sites within the survey area would not have been located. Indeed, we found that regional differences in wild colony density were no longer significant after controlling for both climate and survey area.

The effect of survey area 5.2

We found that survey area had a strong negative correlation with wild colony density. This might be because a greater proportion of

colonies are not detected when survey efforts are spread across larger areas (>50 km²). In contrast, high densities produced by small survey areas $(<5 \text{ km}^2)$ might be the result of a biasing effect whereby surveys are made in small areas where wild colonies are known to be abundant and do not reflect the density of wild colonies across the wider area in which the colonies forage. For instance, Oldroyd et al. (1994) located 37 wild colonies in a small area (0.25 km²) of riparian woodland in Victoria, Australia and reported a density of 148/ km². However, the density would have been considerably lower if the survey area had been extended to include neighbouring habitats that were unsuitable for nesting (i.e. without trees), but in which the wild colonies were foraging. If the foraging area is taken as a circle of radius 2, 3 or 5 km then the actual area would be approximately 12.6, 28.3 and 78.5 km², leading to densities of approximately 2.9, 1.3 and 0.5/km². Of course, there may well have been additional wild colonies in these wider areas, so it is not possible to determine the actual densities. However, it is clear that colony density in a nest site aggregation provided by a restricted area of suitable nesting habitat can be much higher than a colony density relevant to the foraging area of those colonies. Similar aggregations can also occur in Apis dorsata, a species that has open nests, where many colonies nest close to each other under branches of a tree or on a cliff (Oldroyd et al., 2000).

5.3 | The effect of climate and land use

Temperature and net primary productivity were both positively correlated with wild colony density, which probably reflects an increase in foraging activity and the temporal availability of floral resources. Wild A. *mellifera* colonies exhibit seasonal migration in tropical Africa and America, which is considered an adaptation to changes in the spatial distribution of floral resources as it allows them to forage throughout much of the year (McNally & Schneider, 1992). In regions with lower mean annual temperatures, colonies must survive longer winters during which floral resources are scarce or absent and when it is often too cold to forage. In these regions, wild colonies experience elevated mortality during the winter months (up to 80% of founder colonies; Seeley, 2017) and this likely has a large effect on their density the following spring.

Wild colony densities peaked at mean annual temperatures of 23°C (Figure 4), which is consistent with an optimum foraging temperature of approximately 20°C (Abou-Shaara et al., 2017). At mean annual temperatures exceeding 25°C, honey bee colonies are exposed to temperatures that negatively affect foraging and other aspects of colony productivity (Abou-Shaara et al., 2017). Under these conditions, wild colony densities are probably limited by rainfall, which has been shown to be an important factor in seasonally arid locations (Baum et al., 2008; Loper et al., 2006; Oldroyd et al., 1994). Here, we show that rainfall is a less important factor on a global scale, which is consistent with Jaffé et al. (2010) who found that wild colony densities correlated with temperature, but not precipitation, at 25 sites across *A. mellifera*'s native range.

WILEY

Land use and net primary productivity (an index of vegetation) did not significantly affect density at the 41 sampled locations. Honey bees are generalists and can forage over long distances in a variety of habitats (Beekman & Ratnieks, 2000; Ricigliano et al., 2019; Samuelson et al., 2020) so it is likely that land use only affects wild colony density at small spatial scales. In our analysis, land use was generalised over large areas, so we were unlikely to detect smallscale variation in wild colony density, which is arguably more ecologically relevant (Utaipanon, Holmes, et al., 2019).

5.4 | Comparisons with FAO data on managed colonies

Approximately half of all managed honey bee colonies worldwide (45.3 million) are in Asia (FAO, 2021) and the majority of these colonies are assumed to be A. mellifera of European descent (Osterman et al., 2021). Therefore, it is likely that managed swarms frequently escape into the wild, but it seems that they are unable to form self-sustaining wild populations, although this needs to be verified. Proposed explanations include competition with native honey bees (e.g. A. cerana; Manila-Fajardo & Cleofas, 2003), effects of native honey bee parasites (e.g. Tropilaelaps clareae; Oldroyd & Nanork, 2009) and difficulty in regulating brood production in tropical regions with little variation in day length (Rinderer, 1988). Indeed, European honey bees are poorly adapted to tropical climates (Harrison & Hall, 1993) and did not establish large wild populations in tropical America before Africanised honey bees were introduced (Michener, 1975; Quezada-Euán et al., 1996).

Europe has the second-highest number of managed honey bee colonies worldwide at 25.4 million (FAO, 2021). This includes data from 33 countries with a total landmass of 21.7 km² and equates to a density of 1.2 managed colonies/km² which is over four times higher than the average wild colony density reported in Europe. This suggests that a smaller proportion of colonies are subject to natural selection in Europe and that beekeeper management plays a more prominent role in the survival of both managed and wild colonies. For instance, the widespread use of veterinary treatments by beekeepers in Europe might help keep levels of pests and disease low enough for both managed and wild colonies to survive (Thompson, 2012). However, there are probably hotspots in Europe where wild colonies outnumber managed (Requier et al., 2020), and there has been an increased emphasis on natural beekeeping in recent years (Neumann & Blacquiere, 2016) where, amongst other suggestions, beekeepers are encouraged not to treat their colonies with chemicals so that they can evolve a natural resistance to disease (Seeley, 2019). Therefore, natural selection probably still contributes to colony survival in Europe, but not to the same extent as in other regions like Africa where wild colonies are more numerous and commercial beekeeping is poorly developed (Dietemann et al., 2009; Gratzer et al., 2021).

5.5 | Estimated number of wild A. mellifera colonies

Based on the mean regional densities in Figure 3 and the estimated area of habitable landmass in each region, we estimate that there are approximately 280 million wild *A. mellifera* colonies worldwide, which is more than double the number of managed colonies reported by the FAO in 2021 (102 million). This is a tentative estimate based on limited data and should be used with caution. However, the strong indication is that wild *A. mellifera* colonies outnumber managed colonies in most regions, with the exception of Europe.

5.6 | Areas for future research

Our study reveals several knowledge gaps regarding wild A. mellifera colony density which could be addressed by future research. An important area for future research concerns the limits to wild A. mellifera's geographical range. In Europe, wild colonies are thought to occur as far north as 60 degrees latitude (Figure 2), which is consistent with the northernmost survey in our sample (Seeley & Chilcott, 2020). However, permanent beekeeping is practiced as far north as 68 degrees in Finland (Meyer-Rochow, 2008) and it is possible that escaped swarms occur in northern settlements throughout the summer, but whether they survive the long winters at these latitudes is unknown. Similarly, little is known about wild colonies in the southernmost parts of A. *mellifera's* range, such as temperate South America. For instance, there is little information regarding wild colonies in the southern half of Argentina (below Buenos Aires), where the majority are of European descent (Rinderer et al., 1993). This also applies to much of Asia, where wild A. *mellifera* colonies are thought to be absent (Oldroyd & Nanork, 2009).

Analyses of wild colony density and numbers on smaller spatial scales, and possibly incorporating the effects of land use, might help identify hotspots in native regions where wild colonies outnumber managed (Requier et al., 2020). This would have implications for the conservation of native subspecies, given that wild colonies in these areas might represent local ecotypes and an important source of genetic diversity (Requier et al., 2019). For instance, wild colonies in Ireland are considered to be pure *A. mellifera mellifera* (Browne et al., 2020; Hasset et al., 2018), the subspecies native to Northern Europe. Although, managed colonies in Ireland are also mainly *A. mellifera mellifera* (NIHBS, 2021). The degree to which wild colonies are genetically distinct or significantly more native than managed colonies in other parts of Europe is an important topic for future research but is beyond the scope of this review.

6 | CONCLUSION

In French, the honey bee (A. *mellifera*) is called 'l'abeille domestique', the domestic bee and in many countries, presumably including France, it is seen primarily as a bee that lives under human management in hives. However, our study clearly shows that the honey bee

is also 'une abeille sauvage', that is living wild in unmanaged colonies (Seeley, 2019). Indeed, our results indicate that wild colonies outnumber managed colonies, although not in Europe. The realisation of this important fact should have significant consequences on how we view the honey bee. For example, the vast numbers of wild colonies, not to mention the approximately 102 million managed colonies, surely mean that the word endangered, which is frequently used by the media in the context of the honey bee, is far from accurate even though beekeepers, especially those in North America and Europe, have faced increased challenges in maintaining healthy live colonies in recent decades (Genersch, 2010). On a positive note, it also shows that in surviving its challenges the honey bee will be aided by natural selection on wild colonies in many locations. This is shown, for example, by wild colony surveys in New York States's Arnot Forest several decades apart using the bee-lining method (Seeley, 2007; Visscher & Seeley, 1982), which showed the same colony density before and after the arrival of Varroa mites which are now found in the wild colonies (Seelev. 2019).

AUTHOR CONTRIBUTIONS

Oliver D. Visick: Conceptualization (equal); data curation (lead); formal analysis (lead); visualization (lead); writing – original draft (lead). Francis L. W. Ratnieks: Conceptualization (equal); funding acquisition (lead); supervision (lead); writing – original draft (supporting); writing – review and editing (lead).

ACKNOWLEDGEMENTS

The authors are grateful for the comments from the editor and three anonymous reviewers which helped improve the manuscript.

FUNDING INFORMATION

This work was supported by the C.B. Dennis British Beekeepers' Research Trust.

CONFLICT OF INTEREST STATEMENT

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

DATA AVAILABILITY STATEMENT

The data that supports the findings of this study are available in the Supporting Information of this article. Data including R script and QGIS Shapefiles will also be uploaded to Dryad upon acceptance.

ORCID

Oliver D. Visick (D) https://orcid.org/0000-0001-8980-0482

REFERENCES

- Abou-Shaara, H. F., Owayss, A. A., Ibrahim, Y. Y., & Basuny, N. K. (2017). A review of impacts of temperature and relative humidity on various activities of honey bees. *Insectes Sociaux*, 64, 455–463.
- Arundel, J., Oldroyd, B. P., & Winter, S. (2012). Modelling honey bee queen mating as a measure of feral colony density. *Ecological Modelling*, 247, 48–57.

Ecology and Evolution

- Arundel, J., Oldroyd, B. P., & Winter, S. (2013). Modelling estimates of honey bee (*Apis* spp.) colony density from drones. *Ecological Modelling*, 267, 1–10.
- Arundel, J., Oxley, P. R., Faiz, A., Crawford, J., Winter, S., & Oldroyd, B. P. (2014). Remarkable uniformity in the densities of feral honey bee, *Apis mellifera Linnaeus*, 1758 (Hymenoptera: Apidae) colonies in south eastern Australia. *Austral Entomology*, *53*, 328–336.
- Baum, K. A. (2003). Feral Africanized honey bee ecology in a coastal prairie landscape (PhD thesis). Texas A&M University.
- Baum, K. A., Rubink, W. L., Pinto, M. A., & Coulson, R. N. (2005). Spatial and temporal distribution and nest site characteristics of feral honey bee (Hymenoptera: Apidae) colonies in a coastal prairie landscape. Environmental Entomology, 34, 610–618.
- Baum, K. A., Tchakerian, M. D., Thoenes, S. C., & Coulson, R. N. (2008). Africanized honey bees in urban environments: A spatio-temporal analysis. *Landscape and Urban Planning*, 85, 123–132.
- Beekman, M., & Ratnieks, F. L. W. (2000). Long-range foraging by the honey-bee, Apis mellifera L. Functional Ecology, 14, 490–496.
- Bigio, G., Al Toufailia, H., Hughes, W. O. H., & Ratnieks, F. L. W. (2014). The effect of one generation of controlled mating on the expression of hygienic behaviour in honey bees. *Journal of Apicultural Research*, 53, 563–568.
- Bila Dubaić, J., Simonović, S., Plećaš, M., Stanisavljević, L., Davidović, S., Tanasković, M., & Ćetković, A. (2021). Unprecedented density and persistence of feral honey bees in urban environments of a large SE-European city (Belgrade, Serbia). *Insects*, 12, 1127.
- Boreham, M. M., & Roubik, D. W. (1987). Population change and control of Africanized honey bees (Hymenoptera: Apidae) in the Panama Canal area. Bulletin of the Entomological Society of America, 33, 34–39.
- Browne, K. A., Hassett, J., Geary, M., Moore, E., Henriques, D., Soland-Reckeweg, G., Ferrari, R., Loughlin, E. M., O'Brien, E., O'Driscoll, S., Young, P., Pinto, M. A., & McCormack, G. P. (2020). Investigation of free-living honey bee colonies in Ireland. *Journal of Apicultural Research*, 60, 229–240.
- Büchler, R., Berg, S., & Le Conte, Y. (2010). Breeding for resistance to Varroa destructor in Europe. Apidologie, 41, 393–408.
- Carreck, N. L. (2008). Are honey bees (Apis mellifera L.) native to the British Isles? Journal of Apicultural Research, 47, 318–322.
- Chorley, G. P. H. (1981). The agricultural revolution in northern Europe, 1750-1880: Nitrogen, legumes, and crop productivity. *The Economic History Review*, 34, 71–93.
- Coppinger, C. R., Ellender, B. R., Stanley, D. A., & Osborne, J. (2019). Insights into the impacts of rural honey hunting in Zambia. African Journal of Ecology, 57, 610–614.
- Couvillon, M. J., Schurch, R., & Ratnieks, F. L. W. (2014). Waggle dance distances as integrative indicators of seasonal foraging challenges. *PLoS One*, 9, e93495.
- Crane, E. (1999). Recent research on the world history of beekeeping. Bee World, 80, 174–186.
- Crittenden, A. N. (2011). The importance of honey consumption in human evolution. *Food and Foodways*, 19, 257–273.
- Cunningham, S. A., Crane, M. J., Evans, M. J., Hingee, K. L., & Lindenmayer, D. B. (2022). Density of invasive western honey bee (*Apis mellifera*) colonies in fragmented woodlands indicates potential for large impacts on native species. *Scientific Reports*, 12, 3603.
- Danka, R. G., Loper, G. M., Williams, J. L., Villa, J. D., Sugden, E. A., Collins, A. M., & Rinderer, T. E. (1994). Abating feral Africanized honey bees (*Apis mellifera* L.) to enhance mating control of European queens. *Apidologie*, 25, 520–529.
- Darchen, R. (1972). Ecologie de quelques trigones (*trigona* sp.) de la Savane de Lamto (Cote d'Ivoire). *Apidologie*, 4, 341-367.
- De la Rúa, P., Jaffé, R., Munoz, I., Serrano, J., Moritz, R. F. A., & Kraus, F. B. (2013). Conserving genetic diversity in the honeybee: Comments on Harpur *et al.* (2012). *Molecular Ecology*, 22, 3208–3210.

- De la Rúa, P., Paxton, R. J., Moritz, R. F. A., Roberts, S., Allen, D. J., Pinto, M. A., Cauia, E., Fontana, P., Kryger, P., Bouga, M., Buechler, R., Costa, C., Crailsheim, K., Meixner, M., Siceanu, A., & Kemp, J. R. (2014). Apis mellifera. The IUCN red list of threatened species, e.T42463639A42463665.
- Delaplane, K. S., & Mayer, D. R. (2000). Crop pollination by bees. Centre for Agriculture and Bioscience International (CABI).
- Dietemann, V., Pirk, C. W. W., & Crewe, R. M. (2009). Is there a need for conservation of honeybees in Africa? Apidologie, 40, 285–295.
- Food and Agriculture Organization of the United Nations. (2021). Beehive production. FAOSTAT statistical database. https://www.fao. org/faostat/en/#data/QCL
- Gallai, N., Sallesc, J., Settled, J., & Vaissierea, B. E. (2009). Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecological Economics*, 68, 810–821.
- Galton, D. (1971). Survey of a thousand years of beekeeping in Russia. Bee Research Association.
- Gambino, P., Hoelmer, K., & Daly, H. V. (1990). Nest sites of feral honey bees nests in trees. *Apidologie*, *57*, 35–45.
- Genersch, E. (2010). Honey bee pathology: Current threats to honey bees and beekeeping. Applied Microbiology and Biotechnology, 87, 87–97.
- Goodman, R. D., & Hepworth, G. (2004). Densities of feral honey bee Apis mellifera colonies in Victoria. Victorian Naturalist, 121, 210–214.
- Goulson, D., & Derwent, L. C. (2004). Synergistic interactions between an exotic honeybee and an exotic weed: Pollination of *Lantana camara* in Australia. Weed Research, 44, 195–202.
- Gratzer, K., Wakjira, K., Fielder, S., & Brodschneider, R. (2021). Challenges and perspectives for beekeeping in Ethiopia. A review. Agronomy for Sustainable Development, 41, 46.
- Harris, I., Osborn, T. J., Jones, P., & Lister, D. (2020). Version 4 of the CRU TS monthly high-resolution gridded multivariate climate dataset. *Science Data*, 7, 109.
- Harris, J. W. (2007). Bees with Varroa-sensitive hygiene preferentially remove mite-infested pupae aged < five days post capping. *Journal* of Apicultural Research, 46, 134–139.
- Harrison, J. F., & Hall, H. G. (1993). African-European honeybee hybrids have low nonintermediate metabolic capacities. *Nature*, 363, 258–260.
- Hasset, J., Browne, K. A., McCormack, G. P., Moore, E., Society, N. I. H. B., Soland, G., & Geary, M. (2018). A significant pure population of the dark European honey bee (*Apis mellifera mellifera*) remains in Ireland. *Journal of Apicultural Research*, *57*, 337–350.
- Hawkins, G. P., & Martin, S. J. (2021). Elevated recapping behaviour and reduced *Varroa destructor* reproduction in natural *Varroa*-resistant *Apis mellifera* honey bees from the UK. *Apidologie*, *52*, 647–657.
- Hinson, E. M., Duncan, M., Lim, J., Arundel, J., & Oldroyd, B. P. (2015). The density of feral honey bee (*Apis mellifera*) colonies in South East Australia is greater in undisturbed than in disturbed habitats. *Apidologie*, 46, 403-413.
- Hothorn, T., Bretz, F., & Westfall, P. (2008). Simultaneous inference in general parametric models. *Biometrical Journal*, *50*, 346–363.
- Ilyasov, R. A., Kosarev, M. N., Neal, A., & Yumaguzhin, F. G. (2015). Burzyan wild-hive honeybee A. M. mellifera in South Ural. Bee World, 92, 7-11.
- Jaffé, R., Dietmann, V., Allsopp, M. H., Costa, C., Crewe, R. M., Dall'Olio, R., De la Rúa, P., El-Niweiri, M. A. A., Fries, I., Kezic, N., Meusel, M. S., Paxton, R. J., Shaibi, T., Stolle, E., & Moritz, R. F. A. (2010). Estimación de la densidad de colonias de abejas en su rango natural para cerrar la brecha en los censos del declive de polinizadores. *Conservation Biology*, 24, 583–593.
- Jensen, A. B., Palmer, K. A., Chaline, N., Raine, N. E., Tofilski, A., Martin, S. J., Pedersen, B. V., Boomsma, J. J., & Ratnieks, F. L. W. (2005). Quantifying honey bee mating range and isolation in semi-isolated valleys by DNA microsatellite paternity analysis. *Conservation Genetics*, 6, 527–537.

14 of 15

- Kajobe, R., & Roubik, D. W. (2006). Honey-making bee colony abundance and predation by apes and humans in a Uganda forest reserve. *Biotropica*, 38, 210–218.
- Kerr, W. E. (1971). Contribuicao a ecogenetica de algumas espécies de abelhas. *Ciencia e Cult*, 23, 89–90.
- Klein, A., Vaissiere, B. E., Cane, J. H., Steffan-Dewenter, I., Cunningham, S. A., Kremen, C., & Tscharntke, T. (2007). Importance of pollinators in changing landscapes for world crops. *Proceedings of the Royal Society B*, 274, 303–313.
- Koetz, A. H. (2013). Ecology, behaviour and control of Apis cerana with a focus on relevance to the Australian incursion. Insects, 4, 558–592.
- Kohl, P. L., & Rutschmann, B. (2018). The neglected bee trees: European beech forests as a home for feral honey bee colonies. *PeerJ*, 6, e4602.
- Kohl, P. L., Rutschmann, B., Sikora, L. G., Wimmer, N., Zahner, V., D'Alvise, P., Hasselmann, M., & Steffan-Dewenter, I. (2023). Parasites, depredators, and limited resources as potential drivers of winter mortality of feral honeybee colonies in German forests. *Oecologia*, 202, 465–480.
- Kohl, P. L., Rutschmann, B., & Steffan-Dewenter, I. (2022). Population demography of feral honeybee colonies in central European forests. *Royal Society Open Science*, 9, 220565.
- Le Conte, Y., Vaublanc, G., Crauser, D., Jeanne, F., Rousselle, J. C., & Becard, C. M. (2007). Honey bee colonies that have survived *Varroa destructor*. *Apidologie*, 38, 566–572.
- Loper, G. M., Sammataro, D., Finley, J., & Cole, J. (2006). Feral honey bees in Southern Arizona, 10 years after Varroa infestation. American Bee Journal, 146, 521–524.
- Mader, E., Spivak, M., & Evans, E. (2010). Managing alternative pollinators: A handbook for beekeepers, growers, and conservationists. Sustainable Agriculture Research and Education.
- Manila-Fajardo, A. C., & Cleofas, R. (2003). Performance of honeybees (Apis mellifera L.) in three ecosystems in Laguna, Philippines. Philippine Agricultural Scientist, 86, 146–157.
- McNally, L. C., & Schneider, S. S. (1992). Seasonal cycles of growth, development and movement of the African honey bee, *Apis mellifera scutettata*, in Africa. *Insects Sociaux*, 39, 162–179.
- McNally, L. C., & Schneider, S. S. (1996). Spatial distribution and nesting biology of colonies of the African honey bee *Apis mellifera scutellata* (Hymenoptera: Apidae) in Botswana, Africa. *Environmental Entomology*, 25, 643–652.
- Meyer-Rochow, V. B. (2008). Insects and their uses in a cold country: Finland. *Entomological Research*, *38*, 28–37.
- Michener, C. D. (1975). The Brazilian bee problem. Annual Review of Entomology, 20, 399-416.
- Mondet, F., Beaurepaire, A., McAfee, A., Locke, B., Alaux, C., Blanchard, S., Danka, B., & Le Conte, Y. (2020). Honey bee survival mechanisms against the parasite Varroa destructor: A systematic review of phenotypic and genomic research efforts. International Journal for Parasitology, 50, 433–447.
- Moritz, R. F. A., Dietemann, V., & Crewe, R. M. (2008). Determining colony densities in wild honeybee populations (*Apis mellifera*) with linked microsatellite DNA markers. *Journal of Insect Conservation*, 12, 455–459.
- Moritz, R. F. A., Kraus, F. B., Huth-Schwarz, A., Wolf, S., Carrillo, C. A. C., Paxton, R. J., & Vandame, R. (2013). Number of honey bee colonies in areas with high and low beekeeping activity in Southern Mexico. *Apidologie*, 44, 113–120.
- Moritz, R. F. A., Kraus, F. B., Kryger, P., & Crewe, R. M. (2007). The size of wild honeybee populations (*Apis mellifera*) and its implications for the conservation of honeybees. *Journal of Insect Conservation*, 11, 391–397.
- Morse, R. A., Camazine, S., Ferracane, M., Minacci, P., Nowogrodzki, R., Ratnieks, F. L. W., Spielholz, J., & Underwood, B. A. (1990). The population density of feral colonies of honey bees (Hymenoptera: Apidae) in a city in upstate New York. *Journal of Economic Entomology*, 83, 81–83.

- Native Irish Honey Bee Society. (2021). The native Irish honey bee Apis mellifera mellifera. Native Irish Honey Bee Society (NIHBS).
- Neumann, P., & Blacquiere, T. (2016). The Darwin cure for apiculture? Natural selection and managed honeybee health. *Evolutionary Applications*, 10, 226–230.
- Oddie, M. A. Y., & Dahle, B. (2021). Insights from Norway: Using natural adaptation to breed Varroa-resistant honey bees. Bee World, 98, 38–43.
- Oddie, M. A. Y., Dahle, B., & Neumann, P. (2018). Reduced postcapping period in honey bees surviving *Varroa destructor* by means of natural selection. *Insects*, *9*, 149.
- Oldroyd, B. P. (1998). Controlling feral honey bee, *Apis mellifera* L. (Hymenoptera: Apidae), populations in Australia: Methodologies and costs. *Australian Journal of Entomology*, *37*, 97–100.
- Oldroyd, B. P., Lawler, S. H., & Crozier, R. H. (1994). Do feral honey bees (*Apis mellifera*) and regent parrots (*Polytelis anthopeplus*) compete for nest sites? *Australian Journal of Ecology*, 19, 444–450.
- Oldroyd, B. P., & Nanork, P. (2009). Conservation of Asian honey bees. Apidologie, 40, 296–312.
- Oldroyd, B. P., Osborne, K. E., & Mardan, M. (2000). Colony relatedness in aggregations of *Apis dorsata* Fabricius (Hymenoptera, Apidae). *Insectes Sociaux*, 47, 94–95.
- Oldroyd, B. P., Thexton, E. G., Lawler, S. H., & Crozier, R. H. (1997). Population demography of Australian feral bees (*Apis mellifera*). *Oecologia*, 111, 381-387.
- Oleksa, A., Gawroński, R., & Tofilski, A. (2013). Rural avenues as a refuge for feral honey bee population. *Journal of Insect Conservation*, 17, 465–472.
- Osterman, J., Aizen, M. A., Biesmeijer, J. C., Bosch, J., Howlett, B. G., Inouye, D. W., Jung, C., Martins, D. J., Medel, R., Pauw, A., Seymour, C. L., & Paxton, R. J. (2021). Global trends in the number and diversity of managed pollinator species. *Agriculture, Ecosystems and Environment*, 322, e107653.
- Pacífico, E. C., Efstathion, C. A., Filadelfo, T., Horsburgh, R., Cunha, R. A., Paschotto, F. R., Denes, F. V., Gilardieand, J., & Tellaa, J. L. (2020). Experimental removal of invasive Africanized honey bees increased breeding population size of the endangered Lear's macaw. *Pest Management Science*, *76*, 4141–4149.
- Paini, D. R. (2004). Impact of the introduced honey bee (Apis mellifera) (Hymenoptera: Apidae) on native bees: A review. Austral Ecology, 20, 399-407.
- Panziera, D., Van Langevelde, F., & Blacquière, T. (2017). Varroa-sensitive hygiene contributes to naturally selected Varroa resistance in honey bees. Journal of Apicultural Research, 56, 635–642.
- Paton, D. C. (1996). Overview of feral and managed honeybees in Australia: Distribution, abundance, extent of interactions with native biota, evidence of impacts and future research. Australian Nature Conservation Agency.
- Pérez-Sato, J. A., Châline, N., Martin, S. J., Hughes, W. O. H., & Ratnieks, F. L. W. (2009). Multi-level selection for hygienic behaviour in honeybees. *Heredity*, 102, 609–615.
- Quezada-Euán, J. J. G., Echazarreta, C. M., & Paxton, R. J. (1996). The distribution and range expansion of Africanized honeybees (*Apis mellifera*) in the state of Yucatan, Mexico. Journal of Apicultural Research, 35, 85–95.
- R Core Team. (2023). R: A language and environment for statistical computing. R Foundation for Statistical Computing.
- Rangel, J., Giresi, M., Pinto, M. A., Baum, K. A., Rubink, W. L., Coulson, R. N., & Johnston, J. S. (2016). Africanization of a feral honey bee (*Apis mellifera*) population in South Texas: Does a decade make a difference? *Ecology and Evolution*, *6*, 2158–2169.
- Ratnieks, F. L. W., Piery, M. A., & Cuadriello, I. (1991). The natural nest and nest density of the Africanzed honey bee Chiapas, Mexico. *Canadian Entomologist*, 123, 353–359.
- Requier, F., Garnery, L., Kohl, P. L., Njovu, H. K., Pirk, C. W. W., Crewe, R. M., & Steffan-Dewenter, I. (2019). The conservation of native honey bees is crucial. *Trends in Ecology & Evolution*, 34, 789–798.

- Requier, F., Paillet, Y., Laroche, F., Rutschmann, B., Zhang, J., Lombardi, F., Svoboda, M., & Steffan-Dewenter, I. (2020). Contribution of European forests to safeguard wild honeybee populations. *Conservation Letters*, 13, e12693.
- Ricigliano, V. A., Mott, B. M., Maes, P. W., Floyd, A. S., Fitz, W., Copeland, D. C., Meikle, W. G., & Anderson, K. E. (2019). Honey bee colony performance and health are enhanced by apiary proximity to US Conservation Reserve Program (CRP) lands. *Scientific Reports*, 9, 4894.
- Rinderer, T. E. (1988). Evolutionary aspects of the Africanization of honey-bee populations in the Americas. In G. Needham, R. E. Page, Jr.,
 M. Delfinado-Baker, & C. E. Bowman (Eds.), Africanized honey bees and bee mites (pp. 13–18). Ellis Horwood.
- Rinderer, T. E., Harris, J. W., Hunt, G. J., & de Guzman, L. I. (2010). Breeding for resistance to *Varroa destructor* in North America. *Apidologie*, 41, 409–424.
- Rinderer, T. E., Oldroyd, B. P., & Sheppard, W. S. (1993). Africanised bees in the U.S. *Scientific American*, *269*, 84–90.
- Running, S., & Zhao, M. (2021). MODIS/Terra net primary production gapfilled yearly L4 global 500m SIN grid V061 [data set]. NASA EOSDIS Land Processes DAAC. https://neo.gsfc.nasa.gov/view.php?datas etId=MOD17A2_M_PSN
- Rutschmann, B., Kohl, P. L., Machado, A., & Steffan-Dewenter, I. (2022). Semi-natural habitats promote winter survival of wild-living honeybees in an agricultural landscape. *Biological Conservation*, 266, 109450.
- Samuelson, A. E., Gill, R. J., & Leadbeater, E. (2020). Urbanisation is associated with reduced *Nosema* sp. infection, higher colony strength and higher richness of foraged pollen in honeybees. *Apidologie*, 51, 746–762.
- Saunders, M. E., Goodwin, E. K., Santos, K. C. B. S., Sonter, C. A., & Rader, R. (2021). Cavity occupancy by wild honey bees: Need for evidence of ecological impacts. *Frontiers in Ecology and the Environment*, 19, 349–354.
- Schneider, S., & Blyther, R. (1988). The habitat and nesting biology of the African honey bee Apis mellifera scutellata in the Okavango River Delta, Botswana, Africa. Insectes Sociaux, 35, 167–181.
- Seeley, T. D. (2007). Honey bees of the Arnot Forest: A population of feral colonies persisting with Varroa destructor in the northeastern United States. Apidologie, 38, 19–29.

Seeley, T. D. (2016). Following the wild bees. Princeton University Press.

- Seeley, T. D. (2017). Life-history traits of wild honey bee colonies living in forests around Ithaca, NY, USA. *Apidologie*, 48, 743–754.
- Seeley, T. D. (2019). The lives of bees: The untold story of the honey bee in the wild. Princeton University Press.
- Seeley, T. D., & Chilcott, A. B. (2020). In search of wild colonies of honey bees in the Scottish highlands. *Beecraft*, 6, 8–11.
- Seeley, T. D., & Morse, R. A. (1976). The nest of the honey bee (Apis mellifera L.). Insectes Sociaux, 4, 495–512.
- Seeley, T. D., & Radcliffe, R. W. (2018). Deep forest bee hunting. American Bee Journal, 158, 871–877.
- Seeley, T. D., Tarpy, D. R., Griffin, S. R., Carcione, A., & Delaney, D. A. (2015). A survivor population of wild colonies of European honeybees in the northeastern United States: Investigating its genetic structure. *Apidologie*, 46, 654–666.
- Spottiswoode, C. N., Begg, K. S., & Begg, C. M. (2016). Reciprocal signaling in honeyguide-human mutualism. *Science*, 353, 387–389.
- Taber, S. (1979). A population of feral honey bee colonies. American Bee Journal, 119, 842–847.
- Taylor, M. A., Goodwin, R. M., McBrydie, H. M., & Cox, H. M. (2007). Destroying managed and feral honey bee (Apis mellifera) colonies to eradicate honey bee pests. Crop and Horticultural Science, 35, 313–323.

- Taylor, O. R., & Rowell, G. A. (1988). Drone abundance, queen flight distance, and the neutral mating model for the honey bee, *Apis mellifera*. In G. Needham, R. E. Page, Jr., M. Delfinado-Baker, & C. E. Bowman (Eds.), *Africanized honey bees and bee mites* (pp. 173–183). Ellis Horwood.
- Thompson, C. E. (2012). The health and status of the feral honeybee (Apis mellifera sp) and Apis mellifera mellifera population of the UK, PhD thesis. The University of Leeds.
- Tribe, G., Tautz, J., Sternberg, K., & Cullinan, J. (2017). Firewalls in bee nests—Survival value of propolis walls of wild cape honeybee (*Apis mellifera capensis*). Science of Nature, 104, 29.
- Utaipanon, P., Holmes, M. J., Buchmann, G., & Oldroyd, B. P. (2021). Split or combine? Effects of repeated sampling and data pooling on the estimation of colony numbers obtained from drone genotyping. *Apidologie*, *52*, 620–631.
- Utaipanon, P., Holmes, M. J., Chapman, N. C., & Oldroyd, B. P. (2019). Estimating the density of honey bee (*Apis mellifera*) colonies using trapped drones: Area sampled and drone mating flight distance. *Apidologie*, 50, 578–592.
- Utaipanon, P., Schaerf, T. M., Chapman, N. C., Holmes, M. J., & Oldroyd, B. P. (2021). Using trapped drones to assess the density of honey bee colonies: A simulation and empirical study to evaluate the accuracy of the method. *Ecological Entomology*, *46*, 128–137.
- Utaipanon, P., Schaerf, T. M., & Oldroyd, B. P. (2019). Assessing the density of honey bee colonies at ecosystem scales. *Ecological Entomology*, 44, 291–304.
- van Dooremalen, C., Gerritsen, L., Cornelissen, B., Van der Steen, J. J. M., Van Langevelde, F., & Blacquiere, T. (2012). Winter survival of individual honey bees and honey bee colonies depends on level of *Varroa destructor* infestation. *PLoS One*, *7*, e36285.
- Visscher, P. K., & Seeley, T. D. (1982). Foraging strategy of honeybee colonies in a temperate deciduous forest. *Ecology*, 63, 1790–1801.
- von Frisch, K. (1937). The language of bees. Science Progress, 32, 29-37.
- Wenner, A. M. (1989). "Bee-lining" and ecological research on Santa Cruz Island. American Bee Journal, 129, 808–809.
- Wickham, H. (2016). ggplot2: Elegant graphics for data analysis. Springer Publishing.
- Williamson, E., Groom, S., Utaipanon, P., Oldroyd, B. J., Chapman, N., & Hogendoorn, K. (2022). The reliability of honey bee density estimates from trapped drones. *Apidologie*, 53, 62.
- Wood, B. M., Pontzer, H., Raichlen, D. A., & Marlowe, F. W. (2014). Mutualism and manipulation in Hadza-honeyguide interactions. *Evolution and Human Behavior*, 35, 540–546.
- Youngsteadt, E., Appler, R. H., Lopez-Uribe, M. M., Tarpy, D. R., & Frank, S. D. (2015). Urbanization increases pathogen pressure on feral and managed honey bees. *PLoS One*, 10, e0142031.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Visick, O. D., & Ratnieks, F. L. W. (2023). Density of wild honey bee, *Apis mellifera*, colonies worldwide. *Ecology and Evolution*, 13, e10609. <u>https://doi.org/10.1002/ece3.10609</u>